

Homoclinic snaking in plane Couette flow: bending, skewing, and finite-size effects

J. F. Gibson¹†, and T. M. Schneider²‡,

¹Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824 USA

²Emergent Complexity in Physical Systems Laboratory (ECPS), École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

(Received 25 September 2015; revised 8 January 2016; accepted 10 February 2016)

Invariant solutions of shear flows have recently been extended from spatially periodic solutions in minimal flow units to spatially localized solutions on extended domains. One set of spanwise-localized solutions of plane Couette flow exhibits homoclinic snaking, a process by which steady-state solutions grow additional structure smoothly at their fronts when continued parametrically. Homoclinic snaking is well understood mathematically in the context of the one-dimensional Swift-Hohenberg equation. Consequently, the snaking solutions of plane Couette flow form a promising connection between the largely phenomenological study of laminar-turbulent patterns in viscous shear flows and the mathematically well-developed field of pattern-formation theory. In this paper we present a numerical study of the snaking solutions, generalizing beyond the fixed streamwise wavelength of previous studies. We find a number of new solution features, including bending, skewing, and finite-size effects. We establish the parameter regions over which snaking occurs and show that the finite-size effects of the traveling wave solution are due to a coupling between its fronts and interior that results from its shift-reflect symmetry. A new winding solution of plane Couette flow is derived from a strongly skewed localized equilibrium.

1. Introduction

Invariant solutions of the Navier-Stokes equations are known to play an important role in the dynamics of turbulence at low Reynolds numbers (Kawahara, Uhlmann & van Veen 2012). Invariant solutions in the form of equilibria, traveling waves, and periodic orbits have been computed precisely for canonical shear flows such as pipe flow (Faisst & Eckhardt 2003; Wedin & Kerswell 2004; Duguet, Pringle & Kerswell 2008), plane Couette flow (Nagata 1990; Kawahara & Kida 2001; Viswanath 2007; Gibson, Halcrow & Cvitanović 2009) and plane Poiseuille flow (Waleffe 2001; Gibson & Brand 2014). The development of the invariant-solutions approach to turbulence has largely occurred in the simplified context of small, periodic domains, or “minimal flow units” (Jiménez & Moin 1991). More recently, invariant solutions with localized support have been computed for flows on spatially extended domains. These include spanwise-localized equilibria and traveling waves (Schneider, Marinc & Eckhardt 2010*b*; Schneider, Gibson & Burke 2010*a*; Deguchi, Hall & Walton 2013; Gibson & Brand 2014) in plane Couette flow, and spanwise-localized traveling waves (Gibson & Brand 2014) and a periodic orbit (Zammert & Eckhardt 2014*a*) of plane Poiseuille flow. Avila, Mellibovsky, Roland &

† Email address for correspondence: john.gibson@unh.edu

‡ Email address for correspondence: tobias.schneider@epfl.ch

Hof (2013) computed a streamwise-localized periodic orbit of pipe flow, Mellibovsky & Meseguer (2015) a streamwise-localized periodic orbit of plane Poiseuille flow, Brand & Gibson (2014) a doubly-localized equilibrium solution of plane Couette flow, and Zammert & Eckhardt (2014*b*) a doubly-localized periodic orbit of plane Poiseuille flow. The existence and structure of these spatially localized solutions suggests that they are relevant to large-scale patterns of laminar-turbulent intermittency, such as turbulent stripes, spots, and puffs. For example, the periodic orbit of Avila *et al.* (2013) shares the spatial structure and complexity of turbulent puffs in pipe flow, and its bifurcation sequence provides a compelling explanation of the development of transient turbulence in pipes. The doubly-localized equilibrium of Brand & Gibson (2014) has the characteristic shape and structure of turbulent spots in low-Reynolds plane Couette flow, and for a range of Reynolds numbers sits on the boundary between laminar flow and turbulence. Analysis of these localized solutions has so far focused on their bifurcations from spatially periodic solutions (Chantry, Willis & Kerswell 2014; Mellibovsky & Meseguer 2015) and linear analysis of their decaying tails (Brand & Gibson 2014; Gibson & Brand 2014).

The spanwise-localized invariant solutions of plane Couette flow of Schneider *et al.* (2010*b*), are notable for being the first localized solutions discovered, for their relation to the widely-studied equilibrium solution of Nagata (1990); Clever & Busse (1997); Waleffe (1998) (hereafter NBCW), and for exhibiting the particularly interesting feature of homoclinic snaking. Homoclinic snaking is a process by which the localized solutions grow additional structure at their fronts in a sequence of saddle-node bifurcations when continued parametrically (Woods & Champneys (1999); Burke & Knobloch (2006, 2007*a*); Schneider *et al.* (2010*a*); see also § 3.1). Homoclinic snaking occurs in a number of pattern-forming systems with localized solutions, including binary fluid convection (Batiste & Knobloch 2005) and magneto-convection (Batiste, Knobloch, Alonso & Mercader 2006), and it is well-understood mathematically for the one-dimensional Swift-Hohenberg equation (Burke & Knobloch 2006, 2007*a*; Beck, Knobloch, Lloyd, Sandstede & Wagenknecht 2009). Knobloch (2015) provides a comprehensive review of localization and homoclinic snaking in dissipative systems. Though no explicit connection between the Swift-Hohenberg and the Navier-Stokes equations is known, the striking similarity of the localized plane Couette solutions and the localized solutions of Swift-Hohenberg with cubic-quintic nonlinearity suggests there might be a mathematical connection between the two systems. These similarities include the structure of localization, the snaking behaviour, the even/odd symmetry of the snaking solutions, and the existence of asymmetric rung solutions (Schneider *et al.* 2010*a*). One might envision, for example, that a reduced-order model of the localized solutions (Hall & Sherwin 2010; Hall 2012; Beaume, Chini, Julien & Knobloch 2015) might relate the spanwise variation of their mean streamwise flow to the cubic-quintic Swift-Hohenberg equation. Such a relation would link the mathematically well-developed field of pattern-formation theory to the localized solutions of shear flows cited above, or to recent numerical studies of laminar-turbulent pattern formation in extended shear flows (Barkley & Tuckerman 2005; Duguet, Schlatter & Henningson 2010; Tuckerman, Kreilos, Schrobdsdorff, Schneider & Gibson 2014).

In support of developing such connections between pattern-formation theory and shear flows, we present in this paper a more detailed analysis of the snaking solutions of Schneider *et al.* (2010*b,a*). In particular, we examine the effects of varying the streamwise wavelength L_x of the solutions compared to the fixed $L_x = 4\pi$ of Schneider *et al.* (2010*a*). We find that homoclinic snaking is robust in L_x and that the snaking region moves upwards in Reynolds number with decreasing L_x . The ranges of streamwise wavelength and Reynolds number in which snaking solutions exist is found to be $1.7\pi \leq L_x \leq 4.2\pi$

and $165 \leq Re \leq 2700$. Additionally, we find several interesting solution properties that are suppressed at the parameters studied in Schneider *et al.* (2010a). As L_x decreases below 4π and Re increases above 165, the localized solutions deform appreciably compared to their strictly periodic counterparts, the localized equilibria exhibiting a linear skewing and the traveling waves a quadratic bending. We show that skewing and bending are related to the respective symmetries of the equilibrium and traveling wave solutions, and that bending induces finite-size effects in the traveling waves that scale as the inverse of their spanwise width. In contrast, skewing induces no such finite-size effects on the equilibrium solution. We show that the skewed solutions lead to a new periodically winding form of the NBCW equilibrium solution of plane Couette flow.

The structure of this paper is as follows. § 2 outlines the problem formulation and numerical methods. § 3 describes the features of the localized solutions at fixed streamwise wavelength L_x , including homoclinic snaking, bending, skewing, and finite-size effects. § 4 discusses the effects of varying streamwise wavelength, including the regions of wavelength and Reynolds number over which snaking occurs, the breakdown of snaking outside these regions, and the stability of the solutions. § 5 discusses the periodic pattern in the interior of the localized solutions and its relation to the NBCW solution. The new winding solution is presented in § 5.

2. Problem formulation, methodology, and conventions

Plane Couette flow consists of an incompressible Newtonian fluid between two infinite parallel plates moving at constant relative velocity. The Reynolds number is given by $Re = Uh/\nu$ where U is half the relative wall speed, h is half the distance between the walls, and ν is the kinematic viscosity. The $\mathbf{x} = (x, y, z)$ coordinates are aligned with the streamwise, wall-normal, and spanwise directions, where streamwise is defined as the direction of relative wall motion. After nondimensionalization, the walls at $y = \pm 1$ move at speeds ± 1 in the x direction, and the laminar velocity field is given by $y\mathbf{e}_x$. We decompose the total fluid velocity into a sum of the laminar flow and the deviation from laminar: $\mathbf{u}_{\text{tot}} = y\mathbf{e}_x + \mathbf{u}$. Hereafter we refer to the deviation field $\mathbf{u}(\mathbf{x}, t) = [u, v, w](x, y, z, t)$ as “velocity.” In these terms the laminar solution is specified by $\mathbf{u} = 0$, $p = 0$ and the Navier-Stokes equations take the form

$$\frac{\partial \mathbf{u}}{\partial t} + y \frac{\partial \mathbf{u}}{\partial x} + v\mathbf{e}_x + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + \frac{1}{Re} \nabla^2 \mathbf{u}, \quad \nabla \cdot \mathbf{u} = 0. \quad (2.1)$$

The computational domain $\Omega = [-L_x/2, L_x/2] \times [-1, 1] \times [-L_z/2, L_z/2]$ has periodic boundary conditions in x and z and no-slip conditions at the walls. For spanwise-localized solutions, L_z is set to be large so that Ω approximates a spanwise-infinite domain. We use \hat{L}_z to denote the spanwise wavelength of nearly periodic, small-wavelength patterns within the spanwise-localized solutions; typically $\hat{L}_z \ll L_z$. In the present work we impose zero mean pressure gradient in all computations, leaving the mean (bulk) flow to vary dynamically. As described in Gibson, Halcrow & Cvitanović (2008); Gibson *et al.* (2009), direct numerical simulations are performed with Fourier-Chebyshev spatial discretization with 2/3-style dealiasing and semi-implicit time-stepping. Traveling-wave and equilibrium solutions of (2.1) are computed with a Newton-Krylov-hookstep algorithm. Spatial discretizations are chosen so that truncated Fourier coefficients are $O(10^{-5})$ at most and truncated Chebyshev coefficients are $O(10^{-9})$. The computational domain length L_z is chosen so that the magnitude of velocity is no more than 10^{-3} at $z = \pm L_z/2$. The adequacy of spatial resolution was tested by recomputing solutions at higher resolution. Most of the results discussed in § 3, for example, were computed with $24 \times 33 \times 512$

gridpoints on a $3\pi \times 2 \times 24\pi$ domain, for Reynolds numbers in the range $200 \leq Re \leq 350$. All software and solution data are available for download at www.channelflow.org.

The equilibrium and traveling-wave solutions discussed here are all steady states (in a fixed or traveling frame of reference, respectively), so the energy dissipation rate balances the power input from wall shear instantaneously:

$$D = I = \frac{1}{2L_x} \int_{-L_x/2}^{L_x/2} \int_{-L_z/2}^{L_z/2} \left(\left. \frac{\partial u}{\partial y} \right|_{y=-1} + \left. \frac{\partial u}{\partial y} \right|_{y=1} \right) dx dz. \quad (2.2)$$

Note that D is defined in terms of the deviation velocity \mathbf{u} and not the total velocity \mathbf{u}_{tot} , so that D measures the excess energy dissipation of spanwise-localized solutions over the laminar flow, which has $D = 0$. Since the internal structure of a spanwise-localized solution stays roughly constant as non-laminar structure grows at its fronts, D serves as a good measure of the width of a solution. The lack of L_z normalization makes the D of a spanwise-localized solution insensitive to the choice of spanwise length for the computational domain in which it is embedded.

For discussing the symmetries of the flow we follow the conventions of Gibson & Brand (2014), here adding the action of symmetries on the pressure field. Let

$$\begin{aligned} \sigma_x &: [u, v, w, p](x, y, z) \rightarrow [-u, v, w, p](-x, y, z), \\ \sigma_y &: [u, v, w, p](x, y, z) \rightarrow [u, -v, w, p](x, -y, z), \\ \sigma_z &: [u, v, w, p](x, y, z) \rightarrow [u, v, -w, p](x, y, -z), \\ \tau(\Delta x, \Delta z) &: [u, v, w, p](x, y, z) \rightarrow [u, v, w, p](x + \Delta x, y, z + \Delta z), \end{aligned} \quad (2.3)$$

and let concatenation of subscripts indicate products, e.g. $\sigma_{xy} = \sigma_x \sigma_y$. For (ℓ_x, ℓ_z) -periodic fields we define two half-wavelength translation operators $\tau_x = \tau(\ell_x/2, 0)$ and $\tau_z = \tau(0, \ell_z/2)$. The standard group-theoretic angle-bracket notation indicates the group formed by a set of generators; for example $\langle \sigma_{xy}, \tau_x \sigma_z \rangle = \{e, \sigma_{xy}, \tau_x \sigma_z, \tau_x \sigma_{xyz}\}$, where e is the identity (Dummit & Foote 2004).

3. Solution properties at fixed streamwise wavelength

3.1. Snaking

The primary notable feature of the localized solutions is their *homoclinic snaking*. Under continuation in Reynolds number at fixed streamwise wavenumber, the localized equilibrium and traveling-wave solutions follow curves that snake upwards in the Re, D plane, as shown in detail in figure 1 and over a larger range of D in figure 4(d). Velocity fields corresponding to the labeled points in figure 1(a) are shown in figure 2. Figures 2(a,b,c) show that the traveling-wave solution grows additional structure at the solution fronts as it moves upwards in D along the snaking curve, while the interior structure remains nearly constant. The structure of the fronts is the same at alternating saddle-node points (a,c), while the saddle-node point (b) between them has front structure of opposite streamwise sign. Note that due to the σ_{xy} symmetry of plane Couette flow, every traveling-wave solution \mathbf{u} with wave speed c_x has a symmetric partner $\sigma_{xy} \mathbf{u}$ with wave speed $-c_x$. The $\sigma_{xy} \mathbf{u}$ symmetric partner of figure 2(b) has fronts with the same structure and streamwise sign as figure 2(a,c).

Figure 1(a) also shows ‘‘rung’’ solutions that bifurcate from the equilibrium solution in a pitchfork bifurcation near the saddle-node bifurcation points of the equilibrium and connect to the traveling wave near their saddle-node points (or vice versa). The rung solutions are asymmetric, as if formed from an amplitude envelope that drifts in the




FIGURE 1. (color online) **Homoclinic snaking of localized solutions at $L_x = 3\pi$.** (a) Snaking curves at low D (small spanwise width) for localized equilibrium (EQ), traveling-wave (TW), and rung (RN) solutions. Labels indicate the solutions shown as velocity fields in figure 2. (b) Snaking curves at high D (large spanwise width). Filled circles indicate the points of zero skewing (for equilibria) and zero bending (for traveling waves) along the snaking curves; signed open circles mark the positions and signs of the maxima in magnitude of skewing and bending (see § 3.3). Pressure fields for the labeled points are shown in figure 3. Both subplots are details of the $L_x = 3\pi$ snaking curve shown in figure 4(d).

spanwise direction as the solution grows away from its pitchfork bifurcation point (see Kao, Beaume & Knobloch (2014) fig. 6). The rung solutions can also be understood as a combination of two solutions near the saddle-node bifurcation point, with the same Reynolds number and the same internal structure, but with different widths D . For example, the equilibria marked 2d and 2f in figure 1(a) and depicted as streamwise velocity fields in figure 2(d,f) are indistinguishable within the interior $-5 < z < 5$. But their differing values of D indicate different spanwise widths. The contour lines of the fronts of 2(f) extend towards $|z| \approx 10$, whereas those of 2(d) reach just $|z| \approx 8$. The rung solution shown as figure 2(e) and marked 2e in figure 1(a) can then be understood as splicing together the left half of figure 2(d) and the right half of figure 2(f). This splicing can be done over a range of Re in the interior of the saddle-node bifurcation, i.e. along the black lines of the rung branches shown figure 1(a). The splicing construction is necessarily inexact, since the rung solutions have no symmetries and hence travel in both x and z , compared to the equilibrium, which is fixed. However it is close enough that such spliced velocity fields converge quickly to the rung solutions under Newton-Krylov-hookstep search. The rung solutions in this paper were computed by splicing and refinement, followed by continuation in Reynolds number.

3.2. Symmetries of localized solutions

The differences between traveling waves, equilibria, and rungs are intimately related to the different symmetries of those solutions, which can be understood in terms of symmetry-breaking bifurcations of the more symmetric, spatially periodic NBCW solution. This is discussed in detail in Gibson & Brand (2014); here we present a brief summary. With proper placement of the z origin, the traveling waves have a $\tau_x \sigma_z$ “shift-reflect” symmetry. That is, a traveling-wave solution satisfies $\mathbf{u} = \tau_x \sigma_z \mathbf{u}$ or

$$[u, v, w, p](x, y, z) = [u, v, -w, p](x + \ell_x/2, y, -z). \quad (3.1)$$

Solutions with this symmetry can travel in x but not z , since the inversion in z about the origin locks the z phase of the solution, but no such restriction exists for x . For similar reasons, the traveling waves can have nonzero mean streamwise velocity, but




FIGURE 2. (color online) **Velocity fields of localized solutions** illustrated by contours of streamwise velocity in the $y = 0$ midplane, $u(x, 0, z)$ and arrow plots of the streamwise-averaged cross-stream velocity, $[\bar{v}, \bar{w}](y, z)$. Eight contour levels are evenly spaced between ± 0.9 , with negative u in dashed blue lines and positive in solid red. Contour lines for $\bar{u} = 0$ are superimposed on the arrow plots. (a,b,c) show the traveling-wave (TW) solution at the three successive lower saddle-node bifurcation points marked on the snaking curve in figure 1(a). (e,f) show the equilibrium (EQ) solution at $Re = 260$ above and below an upper saddle-node bifurcation, and (d) shows the rung (RN) solution at $Re = 260$, at points labeled in figure 1(a). The solution is shown for $L_x = 3\pi$ and on a subset of the $L_z = 16\pi$ computational domain.

their mean spanwise velocity must be zero. The $\tau_x\sigma_z$ symmetry of the localized traveling waves arises from a subharmonic-in- z bifurcation of the (ℓ_x, ℓ_z) -periodic NBCW solution, which has symmetries $\langle \tau_x\sigma_z, \tau_{xz}\sigma_{xy} \rangle$ when taken in the spatial phase of Waleffe (2003). The subharmonic-in- z bifurcation necessarily breaks the $\tau_{xz}\sigma_{xy}$ symmetry, since this symmetry implies ℓ_z periodicity, as follows. If $\tau_{xz}\sigma_{xy}\mathbf{u} = \mathbf{u}$, then $(\tau_{xz}\sigma_{xy})^2\mathbf{u} = \mathbf{u}$. But a brief calculation shows that $(\tau_{xz}\sigma_{xy})^2 = \tau(0, \ell_z)$. Thus the bifurcated solution loses the $\tau_{xz}\sigma_{xy}$ symmetry of NBCW and retains only $\tau_x\sigma_z$.

The localized equilibrium solution has σ_{xyz} inversion symmetry, satisfying

$$[u, v, w, p](x, y, z) = [-u, -v, -w, p](-x, -y, -z). \quad (3.2)$$

As a result of the inversion of all velocity components about the origin, the spanwise-localized solutions with this symmetry are prevented from traveling in x or z , and the

spatial average of all velocity components is zero. The σ_{xyz} symmetry of the localized equilibrium arises from a similar bifurcation of a phase-shifted NBCW solution. Shifting the NBCW solution by a quarter-wavelength in z , $\mathbf{u} \rightarrow \tau_z^{1/2} \mathbf{u}$, changes each of its symmetries s to the conjugate symmetry $\tau_z^{-1/2} s \tau_z^{1/2}$ (Gibson *et al.* 2009). A brief calculation shows that the conjugated symmetry group of the phase-shifted NBCW solution is $\langle \tau_{xz} \sigma_{xy}, \sigma_{xyz} \rangle$. The $\tau_{xz} \sigma_{xy}$ symmetry implies ℓ_z -periodicity, as before, so the subharmonic-in- z bifurcation breaks the $\tau_{xz} \sigma_{xy}$ symmetry but retains σ_{xyz} .

The symmetries of the traveling-wave and equilibrium solutions and the lack of symmetry in rung solutions are evident in the velocity-field plots shown in figure 2. The z -mirror, x -shift $\tau_x \sigma_z$ traveling-wave symmetry (3.1) is particularly apparent in the fronts of the midplane u contour plots of figure 2(a,b,c), and an even z -mirror symmetry is apparent in the corresponding x -averaged cross-stream $[\bar{v}, \bar{w}](y, z)$ plots. It is also evident from these plots why the traveling-wave solution travels in x . In each of figure 2(a,b,c), both the $u(x, 0, z)$ plots and the $[\bar{v}, \bar{w}](y, z)$ plots show a clear imbalance between the positive/negative streamwise streaks. In comparison, for the equilibrium solutions, the σ_{xyz} symmetry of the equilibrium matches each streamwise streak at negative z with an equal streak at positive z of opposite sign. The rung solution figure 2(e), in contrast, has no symmetry at all. The lack of symmetry in the rungs is due fundamentally to their symmetry-breaking bifurcations from the traveling-wave and equilibrium solutions. It can also be understood physically as a consequence of the formation of rungs via splicing as described in § 3.1, which clearly breaks the σ_{xyz} symmetry of the equilibrium solution (or the $\tau_x \sigma_z$ symmetry if constructed by splicing traveling waves). The complete lack of symmetry in rung solutions means they generally have nonzero wave speeds and nonzero net velocity in both the stream- and spanwise directions.

3.3. Bending, skewing, and finite-size effects

The equilibrium (EQ), traveling-wave (TW), and rung solutions (RN) shown as velocity fields in figure 2 are at low D and thus have small spanwise width. The three different types of solutions appear at first glance to consist of a few copies of the same spanwise-periodic structure placed side-by-side, with fronts on either side that taper to laminar flow. This description, however, is neither entirely accurate nor complete. First of all, the interior structure of the three types of solutions must differ at least slightly because the solution types move at different wave speeds ($c_x = c_z = 0$ for equilibria, $c_x \neq 0, c_z = 0$ for traveling waves, and $c_x \neq 0, c_z \neq 0$ for the rungs). But further differences between the three solutions types become apparent at higher D and greater spanwise width. In this subsection we show that

- the EQs *skew*, displaying a linear tilt in x against z (figure 3a),
- the TWs *bend*, displaying a quadratic curvature in x against z (figure 3b),
- the EQ snaking region has constant bounds in Re (figure 4d),
- the TW snaking region is wider but converges to the EQ's as D^{-1} (figure 4d),
- the TW's streamwise wavespeed decreases to zero as D^{-1} , (figure 4c),
- the EQ's interior structure is periodic and winds in x, z (figure 3a), and
- the TW's interior structure is nonperiodic and slowly modulated in z (figure 3b).

The common thread among these phenomena is the interplay between the fronts and the interior structure. Much of the above can be understood by assuming that the fronts are the determining structures of the solutions, and viewing the other properties as a consequences of the fronts and their orientations, as determined by the solution symmetries.

In this paragraph we present a brief sketch of the interplay between the fronts, symmetries, and solution properties. A fully detailed presentation follows in the remainder of the subsection. For the equilibrium, the odd symmetry and opposite orientation of the fronts about the origin produces a linear x, z skew within the solution's interior. The uniform linear skew allows for periodic structure in the interior that winds linearly in x, z . The winding periodic structure oscillates with D , but is otherwise independent of the solution's overall spanwise width. Consequently, many equilibrium solution properties are independent of the overall spanwise width. In contrast, for the traveling wave, the even z -mirror symmetry and similar orientation of the fronts produces quadratic x, z bending in the interior. This curvature necessarily breaks the periodicity of the solution's interior structure and couples the interior structure and global properties to the spanwise width. The wave speed, bending, snaking region, and interior modulation of the traveling wave all vary according to the relative size of the fronts to the spanwise width, that is, as D^{-1} .

Bending and skewing are most clearly illustrated in terms of the solution pressure fields for the points marked on the snaking curve of figure 1(b). The interior structure of the equilibrium solution in figure 3(a) is oriented along a diagonal line in the x, z plane, whereas that of the traveling wave in figure 3(b) curves upward in x with increasing z . We call the former effect *skewing* and the latter *bending*. Skewing is an x, z -odd phenomenon associated with the σ_{xyz} equilibrium symmetry (3.1), which gives an odd symmetry $p(x, z) = p(-x, -z)$ in the $y = 0$ midplane. Similarly, bending is x, z -even and associated with the $\tau_x\sigma_z$ traveling-wave symmetry (3.2), which gives an even symmetry $p(x, z) = p(x + \ell_x/2, -z)$ in the midplane. We quantify skew or bend by the slope (dx/dz) or curvature (d^2x/dz^2) of an interpolating function that passes through the local minima and maxima of the midplane pressure field. Measured this way, bending and skewing are nearly constant throughout the interior of any given solution, as illustrated by the lines of constant slope or curvature in figure 3.

It is notable that the fronts of equilibrium and traveling-wave solutions are indistinguishable at maximum skew/bend (for example, the right-hand sides near $z \approx 20$ in figure 3a,b) and also at zero skew/bend (figure 3c,d). The fronts on the left-hand sides are determined from the right by symmetry. For the equilibrium, the odd $p(x, z) = p(-x, -z)$ symmetry means the dx/dz slope of the structure has the same sign and magnitude at both the left and right fronts, so that the two fronts can be connected by a uniform periodic structure with constant slope. Importantly, the constant linear slope means the equilibrium solution can exist two steps higher up in D (spanwise width) on the snaking curve, with the same internal winding structure and the same fronts, simply by adding more of the same interior periodic winding structure (or one step by adding half as much and flipping the solution with σ_z). The fact that the equilibrium solution can be extended in spanwise width this way with no change in interior structure thus explains why it snakes in a fixed region of Reynolds numbers, independently of D .

The even $p(x, z) = p(x + \ell_x/2, -z)$ symmetry of the traveling wave, on the other hand, means that the fronts impose a dx/dz slope with opposite signs at either end, so that the line connecting them generally must curve, as in figure 3(b). We observe two features of this curvature in all localized traveling-wave solutions. First, the curvature is constant throughout the solution interior, so that the slope changes uniformly throughout. Thus in marked contrast to the equilibrium, the pattern on the interior of traveling wave is not periodic, but instead changes smoothly throughout. This is apparent in the changing relative streamwise phase of adjacent pressure minima and maxima of the traveling wave in figure 3(b), but also more subtly in the long- z modulation of the pressure field, which is seen most clearly in the change of negative pressure contours from one end of the solution to the other. The positive pressure field has a similar modulation, which is less apparent




FIGURE 3. (color online) **Bending and skewing.** Contour plots of pressure $p(x, 0, z)$ in the $y = 0$ midplane are shown for localized equilibrium (EQ) and traveling-wave (TW) solutions with maximum and zero skewing and bending, corresponding to points marked on figure 1(b). In (a,c), a line of constant slope passes through the pressure minima and maxima, showing uniformity of skew throughout the interior of the equilibrium solution, with (a) showing maximum skewing and (c) zero skewing. In (b,d), a line of constant curvature shows the uniformity of bending for the traveling wave, with (b) showing maximum bending and (d) zero bending. Eight contour levels are evenly spaced between $p = \pm 0.025$, dashed blue for negative p and solid red for positive. The solution is shown for $L_x = 3\pi$ and on a subset of the $L_z = 16\pi$ computational domain.

here but can be highlighted by a different choice of contour levels. Second, the slopes at the fronts vary between fixed bounds, the same bounds as for equilibria. Consequently, as the solution widens upwards along the snaking curve, the curvature decreases, and the interior structure becomes more periodic.

Finite-size effects and D^{-1} scaling. Figure 4(a,b,c) show bending, skewing, and wave speed as a function of D , in comparison to the Re, D snaking in figure 4(d). Several features are notable. First, the solutions snake twice as fast in Re as in skewing, bending,




FIGURE 4. (color online) **Bending (a), skewing (b), and wave speed (c) in comparison to snaking in Reynolds number (d)** for the equilibrium (EQ), traveling-wave (TW), and rung (RN) solutions at $L_x = 3\pi$. Dotted lines show the D^{-1} envelope of wave speed and bending for the traveling wave and the $D = \text{const}$ envelope of equilibrium skewing. In (a) two independent dotted lines of form $R + cD^{-1}$ are shown. The values of R for the two lines were set as the lower and upper bounds of the equilibrium snaking curve ($R = 236$ and $R = 268$), and the values of c chosen to fit the envelope of the traveling-wave snaking curve. Labeled points correspond to pressure fields shown in figure 3. The breakdown of homoclinic snaking for $D \geq 160$ in (d) occurs when the spanwise growth of the localized solution reaches the edges of the $L_z = 24\pi$ computational domain.

or wave speed. This is due to the fact that the points of maximum magnitude in skewing and bending near the upper saddle-nodes in figure 1(b) have opposite sign. Second, the traveling wave's bending and streamwise wave speed curves are nearly identical (figure 4a and c) in all aspects, including position of minima, maxima, and zeros, D^{-1} scaling, and remarkably, magnitude. Sizable discrepancies between bending and wave speed occur only for $D < 40$, when the traveling wave consists of only a few copies of the interior periodic pattern (e.g. figure 2a,b). The nearly identical magnitudes of nondimensionalized bending and wave speed holds only for $L_x = 3\pi$; at other L_x the two quantities are strongly correlated but differ in magnitude by a factor of two or less.

Third, each of the traveling wave's bending, wave speed, and Re snaking plots has a D^{-1} envelope, whereas the corresponding plots for the equilibrium are constant in D . As argued above, the constancy of the equilibrium's behaviour in D is due to the fact that, with linear skew, the solution can be extended in z and thus bumped up to a higher position on the snaking curve at the same Reynolds number simply by adding another copy of the periodic pattern in the interior. Thus the equilibrium snakes between constant bounds in Reynolds number and skewing. For the traveling wave, on the other hand, if we take the slope dx/dz at the fronts as boundary conditions for constant interior




FIGURE 5. **Core, front, tails structure of the localized solutions.** (a) Maximum magnitude of streamwise velocity ($\max_{xy} |u|$) and magnitude of mean streamwise velocity ($|\langle u \rangle_{xy}|$) as a function of z for the traveling wave at a point of maximum bend ($Re = 275, D = 48$). The $|\langle u \rangle_{xy}|$ line is dotted when $\langle u \rangle_{xy}$ is negative. (b) Magnitude of the k_x th streamwise Fourier mode for $k_x = 0, 1, 2$, as measured by the root-mean-square magnitude of $\hat{\mathbf{u}}_{k_x}(y, z)$ over y as a function of z . The computational domain is $3\pi \times 24\pi$.

curvature d^2x/dz^2 over a spanwise width that scales as D , then the curvature must scale as $D^{-1}dx/dz$. Given that the slope of the fronts oscillates between fixed bounds, the bending then must oscillate between bounds that scale as D^{-1} . For large D the curvature thus approaches zero, and the interior of the solution approaches a constant periodic pattern with skewing, bending, and wave speed approaching zero.

At the point of zero bending (figure 3d), the interior structure of the traveling wave is periodic and practically indistinguishable from the structure of the equilibrium at zero skew (figure 3c). These points occur near low- Re saddle-node bifurcations (figure 1b), suggesting that the reason for the close match in the lower bound in Re of the equilibrium and traveling-wave snaking regions is that the two solutions near the lower bifurcation point differ mainly in the orientation of one front. Lastly, the complete lack of symmetry in rung solutions means that they generally travel in z as well as x ; however, the nondimensionalized z wave speeds are on the order of 10^{-5} .

3.4. Core, front, tail structure

The localized solutions are formed from nearly periodic, large-amplitude core structure that taper into small-amplitude, exponentially decaying tails. The near periodicity of the core and the tapering fronts are apparent in figure 2 and figure 3. Figure 5 illustrates the small-amplitude tails as well, through logarithmic plots of velocity magnitude as a function of the spanwise coordinate z . Figure 5(a) shows $|\langle u \rangle_{xy}|(z)$, the magnitude of the xy -average streamwise flow, and $\max_{xy} |u|(z)$, the maximum over x, y of the magnitude of the streamwise flow. Figure 5(b) shows the root-mean-square magnitude over y of several streamwise Fourier modes as a function of z . The central feature of these plots is the dominant $\exp(-\alpha|z|)$ scaling of the tails (where $\alpha = 2\pi/L_x$), consistent with the linear analysis presented in Gibson & Brand (2014). This analysis showed that for large $|z|$, the tails of spanwise-localized, streamwise-periodic solutions are dominated by the $k_x = \pm 1$ streamwise Fourier modes, which take the form $\hat{\mathbf{u}}_{\pm 1}(y) \exp(\pm 2\pi\alpha i(x - c_x t) - \alpha|z|) + c.c.$. The $\exp(-\alpha|z|)$ scaling of the $k_x = 1$ mode is apparent in figure 5(b). The magnitude of the streamwise velocity in the tails ($\max_{xy} |u|$) is dominated by the $k_x = \pm 1$ modes and thus has the same $\exp(-\alpha|z|)$ scaling, as shown in figure 5(a). The $\exp(-2\alpha|z|)$ scaling of the $k_x = 0$ Fourier mode in figure 5(b) results from a resonance between the $k_x = \pm 1$ modes, which, when summed and substituted into the nonlinearity $\mathbf{u} \cdot \nabla \mathbf{u}$, produce an




FIGURE 6. **Net streamwise flow and wave speed of the traveling wave.** The net streamwise flow $\bar{u} = (L_x L_y)^{-1} \int_{xyz} u \, dx \, dy \, dz$ of the traveling wave at $L_x = 3\pi$ varies between roughly fixed bounds, in comparison to the D^{-1} scaling of the wave speed. Wave speed is magnified by a factor of twenty for visibility.

$\exp(-2\alpha|z|)$ forcing term in the $k_x = 0$ momentum equation. The $k_x = 0$ Fourier mode carries the xy -average velocity, so $|\langle u \rangle_{xy}|$ in figure 5(a) has $\exp(-2\alpha|z|)$ scaling. The dominant $k_x = 1$ mode thus produces a small, decaying, but non-zero and constant-sign mean streamwise velocity in the solution tails.

The mean streamwise flow $\langle u \rangle_{xy}(z)$ of the traveling wave has a number of interesting features due to its z -even symmetry, which results from the $\sigma_z \tau_x$ symmetry of the solution \mathbf{u} . For one, $\langle u \rangle_{xy}(z)$ has the same sign in both tails (positive for the solution depicted in figure 5). Additionally, even z symmetry allows for imbalance between positive and negative mean streamwise velocity when summed across the core and front regions. For example, there are three positive $\langle u \rangle_{xy}$ streaks and four negative large-magnitude streaks across the core and initial front of the traveling wave in figure 5, flanked by two lower-magnitude positive streaks. Summing across the core and fronts, and the weak positive tails, gives a net negative streamwise flow across the entire computational domain. Thus we have, with zero pressure gradient conditions, a steady-state solution whose streamwise flow is net negative in the interior, net positive in the tails, and net negative over the whole flow domain. Figure 6 shows how the net streamwise flow $\bar{u} = 1/(L_x L_y) \int_{xyz} u \, dx \, dy \, dz$ varies along the snaking curve (again, the lack of L_z normalization provides for a measure of the deviation from laminar flow that is insensitive to the computational domain). Note that \bar{u} varies between roughly fixed bounds as the spanwise width (D) increases, because it results from an N versus $N + 1$ imbalance of large-magnitude streamwise streaks of opposite sign. This is in contrast to wave speed and bending, which result from balances between the fixed-sized fronts and the increasing core and therefore scale as D^{-1} . For the localized equilibrium solution \bar{u} is zero and the streamwise flows in the $\pm z$ tails have opposing sign, due to σ_{xyz} symmetry of \mathbf{u} and consequent odd symmetry in $\langle u \rangle_{xy}(z)$.

4. Effects of varying streamwise wavelength

4.1. Snaking region and snaking breakdown

In this section we examine the effects of changing the streamwise wavelength L_x . The central results are that snaking is robust in L_x over the range $1.7\pi \leq L_x \leq 4.2\pi$ or $0.48 \leq \alpha \leq 1.2$, and that the snaking region moves upward in Reynolds number with decreasing L_x , with snaking observed over the range $165 \leq Re \leq 2700$. Each of




FIGURE 7. (color online) **Snaking as a function of streamwise wavelength L_x .** (a) Snaking curves for the equilibrium (EQ), traveling-wave (TW), and rung (RN) solutions at $L_x = 2\pi, 2.5\pi, 3\pi$, and 4π . (b) Detail of snaking curve for $L_x = 2.5\pi$. (c) Snaking breakdown for $L_x \geq 4.2\pi$. (d) Snaking breakdown for $L_x \leq 1.7\pi$.

these bounds is reported to two digits accuracy. Thus the localized solutions and the homoclinic snaking behaviour occur over a wide range of Reynolds numbers, including the $Re \approx 300$ to 400 range where Barkley & Tuckerman (2005) and Duguet *et al.* (2010) observed laminar-turbulent patterns in plane Couette flow. Figure 7(a) shows snaking curves for the localized solutions at a variety of streamwise wavelengths. The interlinked snaking structure of the equilibrium, traveling-wave, and rung solutions is preserved under variation in L_x with the following trends. As L_x decreases, the snaking region moves upwards in Re and widens. As in the $L_x = 3\pi$ case, the width in Re of any given equilibrium snaking curve is constant in D , whereas the amplitude of the traveling-wave snaking region decays as D^{-1} . For $L_x = 4\pi$ the excess amplitude of traveling-wave snaking region over the equilibrium is too small to be observed.

Figure 7(c) and (d) illustrate the breakdown of homoclinic snaking outside the range $1.7\pi \leq L_x \leq 4.2\pi$. Above and below these bounds, the solutions are unable to grow additional structure indefinitely at the fronts. Instead of snaking indefinitely, the solution curves turn around and continue to higher Reynolds numbers at roughly constant spanwise width. For L_x values just beyond the given range, the solutions snake a few times before turning around, as illustrated by the $L_x = 1.6\pi$ and 4.5π curves in figure 7(c,d). Figure 7(b) illustrates a peculiar defect in the homoclinic snaking scenario that occurs at the particular value $L_x = 2.5\pi$. At this streamwise wavelength the snaking curve for the traveling wave solution has two distinct branches. The solid curve labeled “TW branch 1” was continued downward from high Reynolds numbers and small spanwise widths ($D \approx 7$) where it connects smoothly to the other snaking curves via continuation




FIGURE 8. **Snaking region and maximum skewing as a function of wavelength.** (a) The snaking region in Re as a function of L_x for the localized equilibrium. The shaded region indicates the range of Re within which snaking occurs at a given L_x . (b) Maximum magnitude of skewing of the equilibrium solution as a function of L_x . For both figures, dots mark measured values, and the curves are interpolated.

in L_x . However this solution branch does not snake; rather it turns around in a saddle-node bifurcation and continues back to at least $Re = 2000$ at finite spanwise width. In contrast the dot-dashed curve labeled “TW branch 2” does snake upward in D ; it constitutes the bulk of the $L_x = 2.5\pi$ traveling wave snaking curve shown in figure 7(a). The branch-2 traveling wave was obtained from the endpoint of the rung solution at $Re \approx 360$, $D \approx 22.5$. At fixed $L_x = 2.5\pi$, the two branches remain distinct to $Re = 2000$ at least, though they can be connected by continuation in the two parameters Re , L_x . We have observed similar defects in snaking curves at several other values of L_x (not shown). It is possible that the snaking breakdown observed for $L_x < 1.7\pi$ and $L_x > 4.2\pi$ is of this type. That is, there might be branches of the solution curves for such L_x at higher D that are disconnected from the nonsnaking solution curves pictured in figure 7(c,d). We have confirmed this in a few particular instances; for example, by continuing a $D = 90$ solution from $L_x = 4\pi$ to $L_x = 4.3\pi$, where it forms a isola above and disconnected from a lower- D solution branch.

Figure 8(a) shows the snaking region of the localized equilibrium in Reynolds number as a function of the streamwise wavelength L_x . Snaking also occurs when the solutions are continued in L_x with Re fixed, within the same shaded Re , L_x parameter regions. The boundaries of the snaking region in Re decrease roughly exponentially, with L_x for $L_x \leq 3\pi$. As L_x increases to 4.2π , the lower bound of the snaking region approaches a minimum of roughly $Re = 165$. It is notable that the breakdown of snaking at $L_x \approx 4\pi$ closely coincides with the vanishing of the amplitude of the D^{-1} scaling in the traveling wave’s snaking region, as seen figure 7(a) and (c). Similarly, the magnitudes of bending and skewing decrease with increasing L_x , and at $L_x \approx 4\pi$ are too small to be observed in plots of the velocity or pressure fields. Figure 8(b) shows the magnitude of the oscillation in skewing over the snaking curve as a function of L_x , as measured by the slopes of the lines through pressure minima and maxima, as shown in figure 3(b). It is possible that the breakdown of snaking for $L_x > 4.2\pi$ is related to the disappearance of these effects at $L_x \approx 4\pi$.

The breakdown of snaking for both $L_x < 1.7\pi$ and $L_x > 4.2\pi$ appears to be associated with the development of defects in the interior pattern of the solution. Instead of continuing to snake by growing additional structure at the fronts, the solutions develop defects at the spanwise center and continue to high Reynolds numbers. Figure 9 shows




FIGURE 9. (color online) **A defect in a localized solution after the breakdown of snaking.** The streamwise velocity of the $L_x = 4.5\pi$ localized equilibrium solution is shown for $Re = 175$ and $D = 37$, on the upper branch of the $L_x = 4.5\pi$ curve in figure 7(c). Plotting conventions are the same as in figure 2.




FIGURE 10. (color online) **Instability of the localized equilibrium in relation to width and wavelength.** The number of unstable eigenvalues of the localized equilibrium solution as a function of spanwise width (D), overlaid on the Re, D snaking curve for (a) $L_x = 4\pi$, (b) $L_x = 3\pi$.

the defect in the $L_x = 4.5\pi$ equilibrium solution at $(Re, D) = (175, 37)$, on the upper branch of the $L_x = 4.5\pi$ curve in figure 7(c). Defects in the $L_x < 1.7\pi$ solutions are quite weak and appear as subtle variations in visualizations of the velocity and pressure fields. The breakdown of snaking and development of central defects observed here is quite similar to that observed by Kao *et al.* (2014) for the Swift-Hohenberg equation with heterogeneous forcing (see their fig. 8).

4.2. Stability

Figure 10 shows the number of unstable eigenvalues of the $L_x = 4\pi$ and 3π equilibrium solutions in comparison to their Re, D snaking curves. At $L_x = 3\pi$ the equilibrium has a minimum of two or three unstable eigenvalues at small spanwise width (low D). Thus it is not strictly an edge state of the flow (Skufca, Yorke & Eckhardt 2006; Schneider & Eckhardt 2006). However one of the corresponding unstable eigenfunctions is antisymmetric, making the solution an edge state of the flow when constrained to σ_{xyz} symmetry. In both cases there is a general trend toward more unstable modes as the solution grows wider. The $L_x = 3\pi$, $D \approx 100$ solutions depicted in figure 3(a,c) have $O(20)$ unstable eigenvalues. Superimposed on this general trend is an oscillation in which the number of unstable eigenvalues increases and decreases along the snaking curve. For both cases the local maxima (minima) in the number of unstable modes occur at points of




FIGURE 11. (color online) **Extraction of the interior periodic pattern of a spanwise localized solution.** (a) The spanwise-localized equilibrium at point of zero skew, $L_x = 3\pi$, $D = 47$, and $Re = 237$. Vertical lines at $z = \pm 2.85$ mark one copy of the nearly periodic interior structure. (b) An exact periodic equilibrium obtained by Newton-Krylov refinement of the structure extracted from (a), with $L_x, L_z = 3\pi, 5.7$ and $Re = 237$. (c) The lower-branch NBCW equilibrium at the same parameter values as (b), obtained by continuation. Plotting conventions for (a,b,c) are the same as in figure 2. (d) Bifurcation curves for the interior periodic pattern and the NBCW equilibrium for $L_x = 2\pi, 3\pi$ and 4π and aspect ratios $L_x/L_z = 1.74, 1.65, 1.74$ respectively. The periodic interior pattern and lower-branch NBCW equilibrium shown in (b,c) are marked on the $L_x = 3\pi$ curve with a circle and square respectively. The interior pattern extracted from spanwise-localized equilibrium at $L_x = 2\pi$ and 4π are also marked with circles.

maximum (minimum) skewing magnitude. In other words, strongly skewed solutions are more unstable than solutions with weak or zero skew. The same trends occur at $L_x = 2\pi$, with the smallest-width solution starting with six unstable eigenvalues. The trend towards more instabilities with increasing solution size contrasts with the one-dimensional Swift-Hohenberg equation, for which the number of unstable eigenvalues oscillates between zero and two with each cycle along the snaking curve (Burke & Knobloch 2007*a,b*).

5. The periodic pattern of the core

5.1. Relation of the periodic pattern to the NBCW solution

Schneider *et al.* (2010*b*) and Schneider *et al.* (2010*a*) established that the snaking solutions are closely related to the NBCW equilibria, in that they result from a localizing bifurcation of NBCW and resemble the structure of NBCW in their interior. However, the relationship between the core structure and the NBCW solutions is more complicated than previously supposed. In particular, for $L_x \leq 3\pi$, the interior pattern and the NBCW solution lie on distinct solution curves when continued in Reynolds number, although

these curves can be connected by continuation in the higher-dimensional parameter space L_x, L_z, Re .

To compare the interior pattern to the NBCW solution, we extracted one copy of the interior pattern of the equilibrium at a variety of L_x values, as illustrated for $L_x = 3\pi$ in figure 11. We begin in figure 11(a) with a streamwise-localized equilibrium solution at a point along the snaking curve of zero skew, in order to maximize the spanwise periodicity of the interior pattern. The natural spanwise wavelength $\hat{L}_z = 5.70$ of the interior pattern was determined by finding the zeros of $\langle u \rangle_{xy}(z)$ on either side of $z = 0$. In figure 11(a) these points are marked with vertical lines at $z = \pm 2.85$. The nearly periodic interior pattern was then interpolated onto uniformly spaced grid points for a spanwise periodic computational domain of width $L_z = 5.70$ and refined with a Newton-Krylov search to the exact equilibrium shown in figure 11(b). We performed this operation to find the exact equilibrium solution corresponding to the interior periodic pattern at several streamwise wavelengths in the range $2\pi \leq L_x \leq 4\pi$. In each case the divergence and the Gibbs phenomenon of the interpolated field were small and the Newton-Krylov refinement converged quickly onto an exact equilibrium. The natural aspect ratio of the interior pattern was always found to be in the range $1.65 \leq L_x/\hat{L}_z \leq 1.75$.

Figure 11(d) shows bifurcation diagrams for the NBCW solution and exact equilibrium computed from the interior pattern for $L_x = 2\pi, 3\pi$, and 4π . Since these solutions are spanwise as well as streamwise periodic, we use the conventional measure of energy dissipation and wall shear rate

$$D_{tot} = I_{tot} = \frac{1}{2L_x L_z} \int_{-L_x/2}^{L_x/2} \int_{-L_z/2}^{L_z/2} \left(\left. \frac{\partial u_{tot}}{\partial y} \right|_{y=-1} + \left. \frac{\partial u_{tot}}{\partial y} \right|_{y=1} \right) dx dz. \quad (5.1)$$

The circles mark the Re, D_{tot} positions of the interior-pattern equilibrium solutions computed from localized solutions as described above, and the lines indicate the parametric continuation of these solutions in Re . For $L_x = 4\pi$ it was straightforward to continue the NBCW solution to the same aspect ratio and Reynolds number and confirm that the pattern and the NBCW solution were the same. However, as L_x decreases, the upper portion of the solution curve pinches off at a codimension-2 bifurcation point near $L_x = 3\pi$, $Re = 237$, leaving the interior pattern and NBCW on distinct solution curves. This pinching occurs at an L_x value just below the $L_x = 3\pi$ solution curve shown in figure 11(d). For $L_x = 2\pi$, the solution curves for the interior pattern (dash-dot line marked with a circle) and the NBCW solution (dash-dot, no marker) are distinct.

5.2. A winding solution of plane Couette flow

It is possible to compute a spatially periodic *winding* solution from a skewed localized equilibrium, that is, a solution with fundamental domain size L_x, \hat{L}_z that is strictly periodic in x , but whose z periodicity involves a phase shift in x ,

$$\mathbf{u}(x, y, z + \hat{L}_z) = \mathbf{u}(x - \Delta x, y, z). \quad (5.2)$$

One such solution is illustrated in figure 12. The solution was computed starting with a localized equilibrium with strong skewing, like that shown in figure 3(a). An iterative process of continuation in the computational domain length L_z and adjustment of skewing by continuation in Reynolds number was performed to find a localized equilibrium whose interior pattern divided the computational domain evenly ($L_z/\hat{L}_z \doteq N$) and whose skew precisely aligned with the x, z diagonal of the computational domain. N copies of this winding interior pattern were then interpolated onto the computational domain as shown in figure 12 and refined with Newton-Krylov search. The resulting winding




FIGURE 12. (color online) **A winding equilibrium solution of plane Couette flow** at $Re = 268$. The solution is strictly periodic in x but has z periodicity involving a phase shift in x of the form $\mathbf{u}(x, y, z + 5.642) = \mathbf{u}(x - 0.496, y, z)$. Nineteen copies of the $3\pi \times 5.642$ pattern fit in the $3\pi \times 107.2$ periodic computational domain. Contours of the midplane pressure field $p(x, z)$ are shown with the same plotting conventions as figure 3.

solution shown in figure 12 at $Re = 268$ has a $L_x = 3\pi, L_z = 107.2$ computational domain and $N = 19$, giving $\hat{L}_z = L_z/N \doteq 5.462$, $\Delta x = L_x/N \doteq 0.496$, and winding symmetry $\mathbf{u}(x, y, z + 5.642) = \mathbf{u}(x - 0.496, y, z)$. The winding symmetry and $\mathbf{u} = \sigma_{xyz}\mathbf{u}$ were enforced during the Newton-Krylov search. It is likely that other winding solutions of shear flows could be computed without recourse to skewed localized solutions, simply by applying skewing coordinate transformations to known spatially periodic solutions and refining with Newton-Krylov search.

6. Conclusions

We have shown that homoclinic snaking is robust under changes in streamwise wavelength for the spanwise-localized solutions of plane Couette flow of Schneider *et al.* (2010a). Homoclinic snaking occurs for these solutions over streamwise wavelengths in the range $1.7\pi \leq L_x \leq 4.2\pi$ and Reynolds numbers $165 \leq Re \leq 2700$, and the snaking region moves upwards in Re as L_x decreases. The localized equilibrium, traveling-wave, and rung solutions thus exist at arbitrarily large spanwise widths over a wide range of Reynolds numbers. Several new properties of the solutions become apparent as L_x decreases below 4π , most importantly the linear skewing of the equilibrium and quadratic bending of the traveling wave. The traveling wave exhibits finite-size effects such as D^{-1} scaling of the bending, wavespeed, and snaking region, due to the nonuniform structure in the solution core induced by the quadratic bending. The linear skewing of the localized equilibrium solution, on the other hand, induces no such finite-size effects. Its core region is very nearly periodic, close enough that a strictly periodic winding solution can be easily developed from it. The number of instabilities of the localized solutions increase with Reynolds number, with spanwise width, and with skewing. Thus, at a fixed Reynolds number, from a statistical viewpoint one would expect narrow patches of the localized pattern to appear more frequently than wide patches, and with weak rather than strong skewing.

The homoclinic snaking of these localized solutions suggests the Navier-Stokes equations might be related to the Swift-Hohenberg equation under plane Couette flow conditions and for certain parameter ranges and flow states. A primary motivation for this paper is to clarify the parameter ranges and solution structures for which this connection might occur. Our results indicate that homoclinic snaking is a finite-Reynolds, finite-wavelength effect, and that the streamwise wavelength and the Reynolds-number snaking region are strongly coupled. Thus it is unlikely that an analytic understanding of

homoclinic snaking in shear flows will be found via asymptotic analysis in large-Reynolds or large-wavelength limits. If the spanwise-localized solutions are to be understood as a long-wavelength modulation of a small-wavelength, spanwise-periodic pattern, our results show that the periodic pattern is a form of the NBCW solution at aspect ratio $L_x/L_z \approx 1.7$. For $L_x < 3\pi$, under continuation in Reynolds number, the periodic pattern lies on a solution curve distinct from the widely-studied NBCW lower-branch solution.

We see no clear connection between the skewing of the localized equilibrium solution and the skewed laminar-turbulent patterns observed in plane Couette flow by Barkley & Tuckerman (2005, 2007), and Duguet *et al.* (2010). The localized solutions exist over a much wider range of Reynolds numbers than the $300 \leq Re \leq 400$ range of observed laminar-nonlaminar patterns. As shown in figure 7(a), this range depends very much on the streamwise length L_x . For the $L_x = 3\pi$ length studied in § 3, the Reynolds number range for snaking is very similar to $300 \leq Re \leq 400$.

Also, the skewing of the localized equilibrium is a streamwise phase shift of the interior pattern as function of spanwise coordinate, whereas the skewing of observed laminar-turbulent patterns is in the orientation of the boundary between the turbulent patches and the surrounding laminar flow. In contrast, the localized solutions studied here are strictly streamwise periodic, and thus have laminar-nonlaminar boundaries aligned with the streamwise direction. One potential route to finding invariant solutions with skewed laminar-nonlaminar boundaries would be to find a streamwise-localized form of a winding solution like that described in § 5.2. However, the angles observed in the literature for laminar-turbulent patterns are quite different than those observed here. These are typically reported as the angle β between the laminar-turbulent boundary and the streamwise x direction; β and skew dx/dz are related by $\beta = \cot^{-1}(dx/dz)$. Duguet *et al.* (2010) report a range of $20^\circ \leq \beta \leq 70^\circ$ for the boundaries of turbulent patches in plane Couette flow for $324 \leq Re \leq 380$. Barkley & Tuckerman (2005) find the range $15^\circ \leq \beta \leq 65^\circ$ for laminar-turbulent patterns in a tilted minimal flow unit at $Re = 350$. From figure 8, the localized equilibria that snake within this range of Reynolds numbers have skew $0 \leq |dx/dz| \leq 0.15$ or $81 \leq \beta \leq 90$.

Acknowledgments: The authors thank John Burke and Edgar Knobloch for illuminating conversations, and John Burke for assistance in the numerical calculations. We are grateful to reviewer 1 for insightful comments and for pointing out several interesting connections to previous literature. TMS was supported by the Swiss National Science Foundation under grant number 200021-160088. Computations were performed on Trillian, a Cray XE6m-200 supercomputer at the University of New Hampshire supported by the Major Research Instrumentation program of the United States National Science Foundation under grant PHY-1229408.

REFERENCES

- AVILA, M., MELLIBOVSKY, F., ROLAND, N. & HOF, B. 2013 Streamwise-localized solutions at the onset of turbulence in pipe flow. *Phys. Rev. Lett.* **110**, 224502.
- BARKLEY, D. & TUCKERMAN, L. S. 2005 Computational study of turbulent laminar patterns in Couette flow. *Phys. Rev. Lett.* **94**, 014502.
- BARKLEY, D. & TUCKERMAN, L. S. 2007 Mean flow of turbulent-laminar patterns in plane Couette flow. *J. Fluid Mech.* **576**, 1009–137.
- BATISTE, O & KNOBLOCH, E 2005 Simulations of localized states of stationary convection in He-3-He-4 mixtures. *Phys. Rev. Lett.* **95** (24), 244501.
- BATISTE, ORIOL, KNOBLOCH, EDGAR, ALONSO, ARANTXA & MERCADER, ISABEL 2006 Spatially localized binary-fluid convection. *J. Fluid Mech* **560**, 149–158.

- BEAUME, CÉDRIC, CHINI, GREGORY P, JULIEN, KEITH & KNOBLOCH, EDGAR 2015 Reduced description of exact coherent states in parallel shear flows. *Phys. Rev. E* **91** (4), 043010.
- BECK, M., KNOBLOCH, J., LLOYD, D.J.B., SANDSTEDTE, B. & WAGENKNECHT, T 2009 Snakes, ladders, and isolas of localized patterns. *SIAM J. Math. Anal.* **41**, 936–972.
- BRAND, E. & GIBSON, J. F. 2014 A doubly-localized equilibrium solution of plane Couette flow. *J. Fluid Mech.* **750** (R3), 1–12.
- BURKE, J. & KNOBLOCH, E. 2006 Localized states in the generalized swift-hohenberg equation. *Phys. Rev. E* **73**, 056211.
- BURKE, J. & KNOBLOCH, E. 2007a Homoclinic snaking: structure and stability. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **17** (3), 037102.
- BURKE, J. & KNOBLOCH, E. 2007b Snakes and ladders: Localized states in the swift-hohenberg equation. *Phys. Lett. A* **360**, 681–688.
- CHANTRY, MATTHEW, WILLIS, ASHLEY P & KERSWELL, RICH R 2014 Genesis of streamwise-localized solutions from globally periodic traveling waves in pipe flow. *Phys. Rev. Lett.* **112** (16), 164501.
- CLEVER, R.M. & BUSSE, F.H. 1997 Tertiary and quaternary solutions for plane Couette flow. *J. Fluid Mech.* **344**, 137–153.
- DEGUCHI, K., HALL, P. & WALTON, A. 2013 The emergence of localized vortex-wave interaction states in plane Couette flow. *J. Fluid Mech.* **721**, 58–85.
- DUGUET, Y., PRINGLE, C.C.T. & KERSWELL, R.R. 2008 Relative periodic orbits in transitional pipe flow. *Phys. Fluids* **20**, 114102.
- DUGUET, Y., SCHLATTER, P. & HENNINGSON, D.S. 2010 Formation of turbulent patterns near the onset of transition in plane Couette flow. *J. Fluid Mech.* **650**, 119–129.
- DUMMIT, D.S. & FOOTE, R.M. 2004 *Abstract Algebra*. Hoboken, NJ: John Wiley & Sons.
- FAISST, H. & ECKHARDT, B. 2003 Traveling waves in pipe flow. *Phys. Rev. Lett.* **91**, 224502.
- GIBSON, J. F. & BRAND, E. 2014 Spatially localized solutions of planar shear flow. *J. Fluid Mech.* **745**, 25–61.
- GIBSON, J. F., HALCROW, J. & CVITANOVIĆ, P. 2008 Visualizing the geometry of state space in plane Couette flow. *J. Fluid Mech.* **611**, 107–130.
- GIBSON, J. F., HALCROW, J. & CVITANOVIĆ, P. 2009 Equilibrium and traveling-wave solutions of plane Couette flow. *J. Fluid Mech.* **638**, 1–24.
- HALL, PHILIP 2012 Vortex–wave interactions: long-wavelength streaks and spatial localization in natural convection. *J. Fluid Mech.* **703**, 99–110.
- HALL, P. & SHERWIN, S. 2010 Streamwise vortices in shear flows: harbingers of transition and the skeleton of coherent structures. *J. Fluid Mech.* **661**, 178–205.
- JIMÉNEZ, J. & MOIN, P. 1991 The minimal flow unit in near-wall turbulence. *J. Fluid Mech.* **225**, 213–240.
- KAO, H.C., BEAUME, C. & KNOBLOCH, E. 2014 Spatial localization in heterogeneous systems. *Phys. Rev. E* **89**, 012903.
- KAWAHARA, G. & KIDA, S. 2001 Periodic motion embedded in plane Couette turbulence: regeneration cycle and burst. *J. Fluid Mech.* **449**, 291–300.
- KAWAHARA, G., UHLMANN, M. & VAN VEEN, L. 2012 The significance of simple invariant solutions in turbulent flows. *Ann. Rev. Fluid Mech.* **44**, 203–225.
- KNOBLOCH, E. 2015 Spatial localization in dissipative systems. *Ann. Rev. Condens. Matter Phys.* **6**, 325–359.
- MELLIBOVSKY, F. & MESEGUER, A. 2015 A mechanism for localization of nonlinear waves in shear flows. *J. Fluid Mech.* **779** (R3), 1–11.
- NAGATA, M. 1990 Three-dimensional finite-amplitude solutions in plane Couette flow: bifurcation from infinity. *J. Fluid Mech.* **217**, 519–527.
- SCHNEIDER, TOBIAS M & ECKHARDT, BRUNO 2006 Edge of chaos in pipe flow. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **16** (4), 041103.
- SCHNEIDER, T. M., GIBSON, J. F. & BURKE, J. 2010a Snakes and ladders: Localized solutions of plane Couette flow. *Phys. Rev. Lett.* **104**, 104501.
- SCHNEIDER, T. M., MARINC, D. & ECKHARDT, B. 2010b Localised edge states nucleate turbulence in extended plane Couette cells. *J. Fluid Mech.* **646**, 441–451.
- SKUFCA, J.D., YORKE, J.A. & ECKHARDT, B. 2006 The edge of chaos in a parallel shear flow. *Phys. Rev. Lett.* **96**, 174101.

- TUCKERMAN, L.S., KREILOS, T., SCHROBSDORFF, H., SCHNEIDER, T.M. & GIBSON, J.F. 2014 Turbulent-laminar patterns in plane Poiseuille flow. *Phys. Fluids* **26**, 114103.
- VISWANATH, D. 2007 Recurrent motions within plane Couette turbulence. *J. Fluid Mech.* **580**, 339–358.
- WALEFFE, F. 1998 Three-dimensional coherent states in plane shear flows. *Phys. Rev. Lett.* **81**, 4140–4143.
- WALEFFE, F. 2001 Exact coherent structures in channel flow. *J. Fluid Mech.* **435**, 93–102.
- WALEFFE, F. 2003 Homotopy of exact coherent structures in plane shear flows. *Phys. Fluids* **15**, 1517–1534.
- WEDIN, H. & KERSWELL, R. R. 2004 Exact coherent structures in pipe flow: traveling wave solutions. *J. Fluid Mech.* **508**, 333–371.
- WOODS, P.D. & CHAMPNEYS, A.R. 1999 Homoclinic orbits in reversible systems and their applications in mechanics, fluids and optics. *Physica D* **129**, 147–170.
- ZAMMERT, S. & ECKHARDT, B. 2014a Periodically bursting edge states in plane Poiseuille flow. *Fluid Dynamics Research* **46** (4), 041419.
- ZAMMERT, S. & ECKHARDT, B. 2014b Streamwise and doubly-localised periodic orbits in plane Poiseuille flow. *J. Fluid Mech.* **761**, 348–359.